

immediate identification and reduction of the CSF pressure by drainage may be beneficial.

Thanks to our editor, Amy Wirtz Newland.

References

1. Crawford ES, Crawford JL, Safi HJ, et al. Thoracoabdominal aortic aneurysms: preoperative and intraoperative factors determining immediate and long-term results of operations in 605 patients. *J Vasc Surg.* 1986;3:389-404.
2. Svensson LG, Crawford ES, Hess KR, Coselli JS, Safi HJ. Experience with 1509 patients undergoing thoracoabdominal aortic operations. *J Vasc Surg.* 1993;17:357-68 (discussion 368-70).
3. Schepens MA, Defauw JJ, Hamerlijnc RP, De Geest R, Vermeulen FE. Surgical treatment of thoracoabdominal aortic aneurysms by simple crossclamping. Risk factors and late results. *J Thorac Cardiovasc Surg.* 1994;107:134-42.
4. Safi HJ, Campbell MP, Miller CC 3rd, et al. Cerebral spinal fluid drainage and distal aortic perfusion decrease the incidence of neurological deficit: the results of 343 descending and thoracoabdominal aortic aneurysm repairs. *Eur J Vasc Endovasc Surg.* 1997;14:118-24.
5. Estrera AL, Miller CC 3rd, Huynh TT, Porat E, Safi HJ. Neurologic outcome after thoracic and thoracoabdominal aortic aneurysm repair. *Ann Thorac Surg.* 2001;72:1225-30 (discussion 1230-1).
6. Svensson LG, Hess KR, D'Agostino RS, et al. Reduction of neurologic injury after high-risk thoracoabdominal aortic operation. *Ann Thorac Surg.* 1998;66:132-8.
7. Safi HJ, Miller CC 3rd, Estrera AL, et al. Chronic aortic dissection not a risk factor for neurologic deficit in thoracoabdominal aortic aneurysm repair. *Eur J Vasc Endovasc Surg.* 2002;23:244-50.
8. Hill AB, Kalman PG, Johnston KW, Vosu HA. Reversal of delayed onset paraplegia after thoracic aortic surgery with cerebrospinal fluid drainage. *J Vasc Surg.* 1994;20:315-17.
9. Hollier LH, Symmonds JB, Pairorero PC, Cherry KJ, Hallett JW, Gloviczki P. Thoracoabdominal aortic aneurysm repair. Analysis of postoperative morbidity. *Arch Surg.* 1988;123:871-5.
10. Jacobs MJ, de Mol BA, Elenbaas T, et al. Spinal cord blood supply in patients with thoracoabdominal aortic aneurysms. *J Vasc Surg.* 2002;35:30-7.
11. Safi HJ, Miller CC 3rd, Azizzadeh A, Iliopoulos DC. Observations on delayed neurologic deficit after thoracoabdominal aortic aneurysm repair (see comments). *J Vasc Surg.* 1997;26:616-22.
12. Azizzadeh A, Huynh T, Miller CI, Safi H. Reversal of twice-delayed neurologic deficits with cerebrospinal fluid drainage after thoracoabdominal aneurysm repair: a case report and plea for a national database collection. *J Vasc Surg.* 2000;31:592-8.
13. Gravereaux EC, Faries PL, Burks JA, et al. Risk of spinal cord ischemia after endograft repair of thoracic aortic aneurysms. *J Vasc Surg.* 2001;34:997-1003.
14. Hollier LH, Money SR, Naslund TC, et al. Risk of spinal cord dysfunction in patients undergoing thoracoabdominal aortic replacement. *Am J Surg.* 1992;164:210-3 (discussion 213-4).
15. Kouchoukos NT, Masetti P, Rokkas CK, Murphy SF, Blackstone EH. Safety and efficacy of hypothermic cardiopulmonary bypass and circulatory arrest for operations on the descending thoracic and thoracoabdominal aorta. *Ann Thorac Surg.* 2001;72:699-707 (discussion 707-8).
16. Crawford ES, Mizrahi EM, Hess KR, Coselli JS, Safi HJ, Patel VM. The impact of distal aortic perfusion and somatosensory evoked potential monitoring on prevention of paraplegia after aortic aneurysm operation (published erratum appears in *J Thorac Cardiovasc Surg* 1989;97:665). *J Thorac Cardiovasc Surg.* 1988;95:357-67.
17. Crawford ES, Svensson LG, Hess KR, et al. A prospective randomized study of cerebrospinal fluid drainage to prevent paraplegia after high-risk surgery on the thoracoabdominal aorta. *J Vasc Surg.* 1991;13:36-45 (discussion 45-6).
18. Kasirajan K, Dolmatch B, Ouriel K, Clair D. Delayed onset of ascending paralysis after thoracic aortic stent graft deployment. *J Vasc Surg.* 2000;31:196-9.
19. Hutschala D, Fleck T, Czerny M, et al. Endoluminal stent-graft placement in patients with acute aortic dissection type B. *Eur J Cardiothorac Surg.* 2002;21:964-9.
20. Greenberg R, Resch T, Nyman U, et al. Endovascular repair of descending thoracic aortic aneurysms: an early experience with intermediate-term follow-up. *J Vasc Surg.* 2000;31:147-56.
21. Huynh TT, Miller CC 3rd, Safi HJ. Delayed onset of neurologic deficit: significance and management. *Semin Vasc Surg.* 2000;13:340-4.
22. Dake MD, Miller DC, Mitchell RS, Semba CP, Moore KA, Sakai T. The "first generation" of endovascular stent-grafts for patients with aneurysms of the descending thoracic aorta. *J Thorac Cardiovasc Surg.* 1998;116:689-703 (discussion 703-4).
23. Murray MJ, Bower TC, Oliver WC Jr, Werner E, Gloviczki P. Effects of cerebrospinal fluid drainage in patients undergoing thoracic and thoracoabdominal aortic surgery. *J Cardiothorac Vasc Anesth.* 1993;7:266-72.
24. Coselli JS, Lemaire SA, Koksoy C, Schmittling ZC, Curling PE. Cerebrospinal fluid drainage reduces paraplegia after thoracoabdominal aortic aneurysm repair: results of a randomized clinical trial. *J Vasc Surg.* 2002;35:631-9.
25. Safi H, Miller CI, Carr C, Iliopoulos D, Dorsay D, Baldwin J. The importance of intercostal artery reattachment during thoracoabdominal aortic aneurysm repair. *J Vasc Surg.* 1998;27:58-68.

Discussion

Dr Lars G. Svensson (Cleveland, Ohio). I would like to thank the Association for the invitation to comment on the interesting and vexing problem of delayed deficits presented by Dr Safi in a large series of descending and thoracoabdominal aortic repairs. Unfortunately, they did not present the postoperative causes. We have seen it occur with hypotension from things like bleeding, arrhythmia, overdiuresis, sedation, reintubation, and other causes such as pulmonary emboli, arterial embolic showers, thrombosis of the intercostal, or lumbar artery patches or bypasses, and, in some patients, for no apparent reason.

There appear to be at least 3 common pathways: first, loss of spinal cord blood flow, particularly with a tenuous and perilous collateral blood supply; second, failure of adequate spinal cord oxygenation; and third, secondary spinal cord injury from the delayed complex biochemical cascade of deleterious pathways following ischemia and reperfusion.

The preoperative risk factors presented by Dr Safi of type II thoracoabdominal aneurysm, acute dissection, and renal disease reflect the problem of maintaining collateral blood supply in some of these patients. Because you only reattach intercostals from T8 to T12, according to your manuscript, do you think reattaching a wider range of intercostals or lumbar arteries may have reduced the risk? What are you now doing differently? We have found postoperative induced hypertension and delayed extubation useful. Do you think in patients with extensive aneurysms and few intercostal or lumbar artery ostia, endarterectomies should be performed for reattaching segmental arteries?

You had more deficits occur with CSF drainage. Do you think delayed CSF hypertension may be a factor? In a previous prospective randomized study in which postoperative hypotension resulted in 32% of the deficits, CSF drainage tended to be protective. In an excellent study recently published by Coselli, there also appeared to be some possible protection against delayed deficits. In 57% of your patients, delayed deficits improved with repeat drainage. Do you think that delayed removal of the catheter may be useful or do you think this would increase the risk of complications from the catheter too much?

Dr Safi. Thank you, Dr Svensson. With regard to our practice with intercostal arteries, we believe that intercostal arteries 8 to T12 are very important, and our work showed that these intercostal arteries are important for spinal cord perfusion. But in a paper from Amsterdam where they felt that if the lower intercostal arteries are not patent or occluded, the upper intercostal arteries are important, and currently, if we open the aneurysm and the intercostal arteries and the lower ones are not open, then we reattach the T5, 6, and 7. That is the only change.

With regard to the postoperative management, this is an intensivist study by Tam Huynh, and we currently believe that keeping a mean pressure of a 100 mm Hg is the order of the day, because hypotension correlates with paraplegia, as you and I have seen.

With regard to endarterectomy, I caution endarterectomy because I helped my late mentor, Dr Crawford, with 2 cases where the patient developed postoperative bleeding, one 2 hours after the endarterectomy and one 24 hours. I currently don't know how to reattach intercostal arteries to a graft, either an acutely dissected aorta or endarterectomy, because of risk of bleeding with a catastrophic event.

With regard to CSF drainage, we believe that CSF pressure, if it is high, may correlate with paraplegia. How long we are going to keep it, according to our median time for the CSF of the development of neurological deficits, is 2 days, so we are keeping it a day extra. I think if you keep it longer than that there is a risk of infection and some other complication.

Dr Ludwig von Segesser (*Lausanne, Switzerland*). I wish to congratulate Dr Safi for his excellent presentation, and I have a few questions with regard to the extent of repair. As a matter of fact, I did not see the exact proportion of thoracoabdominal repairs versus descending thoracic aortic repairs, and I believe that in acute aortic dissections, most of these cases can be repaired proximally first, and maybe there is a necessity for a distal reentry procedure.

I am asking this because in our environment there is an increasing proportion of endovascular aneurysm repairs, and under these circumstances there is no aortic crossclamping, and of course this may induce a shift or a selection bias in the patients you have referred.

Could you comment on this?

Dr Safi. In our experience, two thirds of our patients are extent I and extent II, so most of them are the more serious aneurysm. With regard to endovascular repair, there are reports in the literature that show delayed paraplegia following the insertion of the stent. With regard to acute dissection, if you remove the upper half of the descending thoracic aorta, risk of death, neurological deficit, or renal failure is minimal, but when we remove the thoracoabdominal aorta completely, from the left subclavian to the iliac bifurcation, the risk in our hands is about 35% to develop a neurologic deficit and the mortality rate is very high.